



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

June 6, 2017

Ms. Stephanie Hacker  
Regulatory Affairs Specialist  
Kelly Registration Systems  
14481 Lochridge Blvd.  
Covington, GA 30014

Subject: Label Amendment – Master Label Update

Product Name: EnProAI Fly Control Block with RABON Oral Larvicide  
Also known as {SWEETLIX Livestock Supplement System; EnProAI RABON Supplement; SWEETLIX RABON Poured Block; and VMS Livestock Mineral-Vitamin Supplement EnProAL Fly Control Block with RABON}

EPA Registration Number: 7698-30  
Application Date: March 10, 2017  
Decision Number: 527680

Dear Ms. Hacker:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false

Page 2 of 2  
EPA Reg. No. 7698-30  
Decision No. 527680

or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Matthew Sellner by phone at 703-347-8034, or via email at [sellner.matthew@epa.gov](mailto:sellner.matthew@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read 'Gebken', written in a cursive style.

Richard Gebken  
Product Manager 10  
Invertebrate & Vertebrate Branch 2  
Registration Division (7505P)  
Office of Pesticide Programs

Enclosure

# EnProAI® Fly Control Block with RABON® Oral Larvicide

{ABNs: SWEETLIX® EnProAI RABON® Supplement;  
SWEETLIX® Livestock Supplement System EnProAI® RABON® Supplement;  
SWEETLIX® RABON® Poured Block; and  
VMS® Livestock Mineral-Vitamin Supplement EnProAI® Fly Control Block with RABON®}

## A Free Choice Supplement Block for Cattle [and Horses]

For control of fecal flies in manure of treated cattle [and horses]. Prevents development of face, horn, house and stable flies in the manure of treated cattle. [Prevents development of house and stable flies in the manure of treated horses.]

**Active Ingredient:**

Tetrachlorvinphos* (CAS # 22248-79-9) .....	0.3%
<b>Other Ingredients**</b> .....	<b>99.7%</b>
<b>Total</b> .....	<b>100.0%</b>

\* RABON® Insecticide - Contains 1.35 grams RABON® per pound

\*\*Refers only to ingredients which are not larvicidal

**Guaranteed Analysis**


*{Will vary according to formulation and state labeling regulations.}*

Crude Protein (Min) .....	7.50%
Crude Fat (Max) .....	0.10%
Crude Fiber (Max) .....	3.00%
Calcium (Min) .....	3.00%
Calcium (Max) .....	4.00%
Phosphorus (Min) .....	1.00%
Salt (Min) .....	9.00%
Salt (Max) .....	10.80%
Magnesium (Min) .....	3.00%
Copper (Min) .....	400 ppm
Iodine (Min) .....	20 ppm
Manganese (Min) .....	1,500 ppm
Selenium (Min) .....	12 ppm
Zinc (Min) .....	1,100 ppm
Vitamin A (Min) .....	120,000 IU/lb
Vitamin D-3 (Min) .....	12,000 IU/lb
Vitamin E (Min) .....	12 IU/lb

**Ingredients**

*{Will vary according to formulation and state labeling regulations.}*

Molasses products, plant protein products, salt, calcium carbonate, magnesium oxide, monocalcium phosphate, dicalcium phosphate, manganous oxide, manganese sulfate, zinc oxide, zinc sulfate, copper sulfate, sodium selenite, ethylenediamine dihydriodide, calcium iodate, cobalt carbonate, Vitamin A supplement, Vitamin D-3 supplement, Vitamin E supplement and mineral oil.



## Keep Out of Reach of Children

### CAUTION

#### First Aid

<b>If Swallowed:</b>	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor for treatment advice.</li><li>• Have person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>If on Skin:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Immediately rinse skin with plenty of water for 15-20 minutes.</li><li>• Call poison control center or doctor for treatment advice.</li></ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first five minutes, then continue rinsing.</li><li>• Call a poison control center or physician for treatment advice.</li></ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For general information on product use, etc., call the National Pesticides Information Center at 1-800-858-7378. For emergencies, call the poison control center at 1-800-222-1222.

#### Note to Physicians and Veterinarians

This product is a cholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. Product may cause aspiration pneumonia. Probable mucosal damage may contraindicate the use of gastric lavage.

*(Note: The First Aid Statements' grid format will be used if market label space permits; otherwise a paragraph format will be used.)*

### PRECAUTIONARY STATEMENTS

**CAUTION:** Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

#### Personal Protective Equipment ("PPE")

Handlers must wear: long-sleeved shirt and long pants; shoes and socks; and chemical-resistant gloves.

#### User Safety Requirements

Follow manufacturers' instructions for cleaning/maintaining PPE. If no such instructions for washables are provided, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### User Safety Recommendations

Users should wash hands with plenty of soap and water before eating, drinking, chewing gum, using tobacco or using the toilet. Users should immediately remove clothing if pesticide gets inside, then wash thoroughly and put on clean clothing. Users should immediately remove PPE after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### Environmental Hazards

This product is highly toxic to fish.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

[Use Restriction: Not to be used on horses destined for slaughter.]

**[Precautions for Horse Use:** This product contains Tetrachlorvinphos, which is a cholinesterase inhibitor. The most frequently reported clinical signs of cholinesterase inhibition in the horse are abdominal pain, lethargy, sweating, tearing and excessive salivation. If these signs are seen in horses, immediately consult your veterinarian. Do not use this product simultaneously or within a week before or after treatment with cholinesterase inhibiting drugs, pesticides or chemicals. Consult a veterinarian before using this product on debilitated, aged, pregnant or nursing animals. This product has not been tested in breeding horses.]

1. Feed 1 block per 5 head of cattle or horses. This allows all animals equal access.
2. Place blocks where animals congregate. Locate blocks throughout pasture, placing them near water and loafing areas. Maintain adequate distance between blocks (10 ft. minimum).
3. Do not allow animals to run out of blocks. When a block is half-consumed, place a new block near it.
4. Feed blocks at the rate of 0.88 oz. per 100 lbs. of bodyweight per day. This intake will supply 0.07 g of larvicide per 100 lbs. of bodyweight per day (2.1 grams per 100 lbs. bodyweight per month). Animals will consume about 8.8 oz per head daily based on a 1,000 lb. animal.
5. Blocks can be fed to animals in confinement. Exact location of blocks to obtain desired consumption will vary between confinement feeding programs.
6. Feed as the only **free choice** source of salt, other minerals and vitamins.
7. **EnProAI Fly Control Block with RABON Oral Larvicide** prevents development of fecal flies in the manure of treated cattle [and horses], but it is not effective against existing adult flies. It is necessary to use other control measures to reduce existing adult fly populations. In order to achieve optimum fly control, use **EnProAI Fly Control Block with RABON Oral Larvicide** in conjunction with other good management and sanitation practices. Start feeding early in the spring before flies begin to appear and continue feeding until cold weather limits fly activity. **EnProAI Fly Control Block with RABON Oral Larvicide** may be fed to cattle up until slaughter and to lactating cows. Follow feeding directions. When animals consume the block at the recommended rate, it will supply 3 mg of selenium. Consumption of selenium should not exceed 3 mg per head daily.

### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place in the original container. **Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Offer for recycling, if available. Dispose of empty container in a sanitary landfill or by incineration, or if allowed by state or local authorities, by burning. If burned, stay out of smoke.

*{Per PR Notice 2007-4, a lot/batch number will appear on either the labeling or container.}*

**IMPORTANT:** This product, which contains added copper, should not be fed to sheep or any species that have a low tolerance to supplemental copper.

For additional information, please contact your {Brand/Company Name} representative.

EPA Reg. No. 7698-30      EPA Est. \_\_\_\_\_  
NET WEIGHT: \_\_\_\_ LBS (\_\_\_\_ KG)

RABON® is a registered trademark of Bayer HealthCare LLC  
EnProAI®, Sweetlix® and VMS® are registered trademarks of Ridley USA Inc.

Manufactured by  
**Ridley USA Inc.**  
111 West Cherry Street, Suite 500  
Mankato, MN 56001

[[Batch #]]  
{Product Code}  
{Date Code}

{Graphics}



{End of Graphics}

[ ] Denotes language that is optional

{ } Denotes language that does not appear on the market labeling.